

5 FirstVillage Drive
P. O. Box 2000
Pinehurst, NC 28374-2000
910-295-6831
www.pinehurstsurgical.com



Surgery Center of Pinehurst MPFL Repair Home Care Instructions

GENERAL & BARIATRIC SURGERY

RAYMOND G. WASHINGTON, M.D.
FABIAN E. ALZAMORA, M.D.
H. WILLY CHU, M.D.
DAVID W. GRANTHAM, M.D.
REID C. VEGELER, M.D.
BAILEY SANDERS, M.D.

EAR, NOSE & THROAT, HEAD & NECK

CARL W. BERK, M.D.
JEFFERSON K. KILPATRICK, M.D.
WYMAN T. MCGUIRT, M.D.
WALDEMAR L. HEFKOHL, M.D.
MATTHEW R. GRAFENBERG, M.D.
JUSTIN D. MILLER, M.D.

WOMEN'S COMPREHENSIVE HEALTH

STEPHEN A. SZABO, M.D.
WILLIAM V. TERRY, M.D.
LISSETTE MACHIN, M.D.
SHAYNA S. JONES, M.D.
TAYLOR MCCAULEY, M.D.
RYAN B. NANCE, D.O.

UROLOGIC SURGERY

ROBERT J. CHAMBERLAIN, M.D.
GREG L. GRIEWE, M.D.
MANLIO A. GOETZL, M.D.
JOHN R. MICHALAK, M.D.

VASCULAR AND VEIN CARE

CLINTON K. ATKINSON, M.D.
PAUL A. VIETA, JR., M.D.
GLENN FUSONIE, M.D.
DAVID ADAMOLEKUN, D.O.

OPHTHALMOLOGY

MILES W. WHITAKER, M.D.

ORTHOPAEDIC SURGERY

MARK E. BRENNER, M.D.
WARD S. OAKLEY, JR., M.D.
NEIL A. CONTI, M.D.
JOHN R. MOORE, IV, M.D.
DAVID E. STROM, M.D.
DAVID J. CASEY, M.D.
KURT P. WOHLRAB, M.D.
DANIEL M. WILLIAMS, M.D.
CHRISTOPHER M. JOHNSON, M.D.
JAMES T. MARINO, M.D.
TONI A. WASHINGTON, M.D.
DANIEL M. WILLIAMS, MD
ALEXANDER C. LEMONS, M.D.
LYNN M. FRASER, M.D.

PLASTIC & FACIAL PLASTIC SURGERY

RUSSELL B. STOKES, M.D.
JEFFERSON K. KILPATRICK, M.D.

Bandages:

Keep dressing clean and dry for 72 hours. All sutures are inside. You may take the dressing off on post-op day #3 and begin showering. Once you take the bandages off there will be steri-strips over the incisions, leave these on until post-operative visit unless they fall off on their own. Keep area clean and dry once bandages are removed.

Activity/Elevation:

Limit your activity for the first 48 hours after surgery. Walking should be limited to what is necessary and comfortable. Elevate your knee with a pillow under your ankle, not your knee for the first 48 hours. This will help to decrease the amount of swelling you may have. Brace should be worn locked in extension when weightbearing, at 0-30 degrees of flexion until post-op visit. Weight bearing on affected leg as tolerated. Use crutches for support when walking.

Ice:

Ice will help relieve the swelling and pain. Use the ice bag on your knee for 48 hours, 20 minutes on and 20 minutes off, during waking hours. If you have been given a **Durakold ice wrap**, please use this as instructed.

Medication:

A prescription for a pain killer will be provided and is usually needed for the first few days after surgery. It may cause nausea and constipation. Please take the medication with food and drink plenty of fluids to stay hydrated. If you experience severe nausea with the medication, please contact my office. Do not drive as long as you are taking pain medication.

Follow-up Appointment:

Please keep your scheduled appointment or call us if it is inconvenient so that we can reschedule you.

Things to watch out for and call the office about are:

Bleeding: if the bandage becomes soaked or the bloody area is larger than the size of a hand, call the office.

Signs of Infection: redness, tenderness, pain, swelling or green/yellow discharge around incision site)

Fever: above 101

Pain: unrelieved by the pain medicine

Difficulty breathing

***If you have any problems or questions after your surgery call**

- **Dr. Wohlrab's Triage Line at 910-621-3791.**
- **After office hours at 910-715-1000 and ask for the Pinehurst Surgical Orthopedic Doctor on call.**
- **Our Afterhours Clinic is also available to walk in Monday-Friday 5PM-7PM, Saturday 9AM-11AM and Sunday 1PM-3PM.**